



AROUND 3 OUT OF 10 PEOPLE with psoriasis will get PsA, so if you're living with psoriasis you may be at risk¹

YOU MAY ALREADY HAVE PsA & not know it – 15% of people with PsA may be undiagnosed²

15%



PsA IS JUST AS DISABLING as rheumatoid arthritis³

For People with PsA:

DAMAGE TO JOINTS & SPINE

Joints can be damaged WITHIN 6 MONTHS of first PsA symptoms⁴



UP TO **47**% MAY HAVE JOINT DAMAGE within two years of initial symptoms⁵

UP TO 40% may have inflammation of the spine, leading to neck & back pain & REDUCED MOBILITY⁶

IMPACT ON EMOTIONAL WELLBEING & OTHER CONDITIONS

Around **20**% experience DEPRESSION⁷

Over **50**% will have another RELATED MEDICAL CONDITION⁸

Conditions related to PsA include:

- cardiovascular disease
- obesity
- diabetes
- Crohn's disease
- eye disease
- depression & anxiety

Daily life with PsA



PAIN & STIFFNESS CAUSED BY PsA CAN BE

DEBILITATING, affecting work, sleep, relationships & quality of life⁹



SIMPLE TASKS CAN ALSO BE AFFECTED, e.g. getting in & out of bed, washing & personal care, dressing, walking & taking part in sports¹⁰

PsA can AFFECT WORK, SLEEP and IMPACT ON QUALITY OF LIFE 9.11



4 OUT OF 10 PEOPLE with PsA describe their disease as A LARGE PROBLEM in everyday life¹²

PsA IS TREATABLE. Consult your doctor if you think you might have PsA.

For more Information: speak to your doctor, visit www.SeePsoriasisLookDeeper.co.uk or contact the Psoriasis Association: www.psoriasis-association.org.uk or call 01604 251620.



References

1. NHS Choices. Psoriatic Arthritis. 2015 Available at: http://www.nhs.uk/conditions/psoriasis/Pages/Introduction.aspx (last accessed October 2015) 2. Villani A.P. Prevalence of undiagnosed psoriatic arthritis among psoriatis patients. Systematic review and meta-analysis. 2015. Journal of the American Academy of Dermatology 3. Sokoll KB, Helliwell PS, "Comparison of disability and quality of life in rheumatoid and psoriatic arthritis.", J Rheumatol., 2001; 28(8): 1842–1846. 4. Haroom M, Gallagher P, FitzGerald O. Diagnostic delay of more than 6 months contributes to poor radiographic and functional outcome in psoriatic arthritis. an early synovitis clinic experience. Rheumatology. 2003;42(12):1460-8. 6. Giadman G, et al. Psoriatic arthritis: a periadio outcome. Ann Rheum Dis 2005; 64(Suppl II): iiit—iiit 7. American College of Rheumatology. Amciety and Depressive Symptoms and Illness Perceptions in Psoriatic Arthritis care 8. Research Vol. 64, No. 10, October 2012; pp. 1593—1601
8. Ogdie A, Schwartzman S, Husni MF, "Recognizing and managing comorbidities in psoriatic arthritis.", Curr Opin Rheumatol, 2015; 27, pp. 118—126
9. Lee. The Burden of Psoriatic arthritis: A permatol, 2014; 70, pp. 671—881
9. It. Tille W, de-Vries C, McHugh M J. "Work disability in psoriatic arthritis: a systematic review", Rheumatology, 2012; 31, pp. 275—283
11. Tille W, de-Vries C, McHugh M J. "Work disability in psoriatic arthritis: a systematic review", Rheumatology, 2012; 31, pp. 275—283
12. Gelfand JM, Giadman DD, Messe P, Let at. Epidemiology of psoriatic arthritis the population of the United States. J Am Acad Dermatol. 2005;53(4):573.